



Hooman Soltanian, M.D., F.A.S.C.

Diplomate American Board of Plastic Surgery

1000 Asylum Ave
Suite 2114
Hartford, CT 06105
Office 860 . 722 . 9544
Fax 860 . 760 . 6363
www.spsect.com

Post-Operative Instructions After Breast Augmentation

- ☞ Take your medications as prescribed.
- ☞ Drink plenty of fluids and try to maintain a high fiber diet.
- ☞ Avoid strenuous activities until you have cleared it with your surgeon.
- ☞ Call your surgeon if you develop a fever higher than 101.5, chills, nausea and vomiting or dizziness.
- ☞ Call your physician if the site of your surgery becomes significantly red and/or swollen.
- ☞ Do not take any Aspirin or Motrin or medication containing aspirin or Motrin for 2 weeks after your surgery, unless instructed by your surgeon.
- ☞ Ambulate several times a day to promote the circulation in your legs.
- ☞ If your calves become painful or you have significant swelling in your legs, please notify our office.
- ☞ Do not drive while taking narcotic pain medication or experiencing significant pain. When you drive you should be able to react to unforeseeable situations without hesitation.
- ☞ Call our office at 860-722-9544 to schedule your follow-up appointment.
- ☞ Keep the surgical bra and the elastic dressing on as much as possible.
- ☞ You may take a shower 48 hours after your surgery.
- ☞ You may wear your own soft sport bra after the second day.
- ☞ You may see some bruising and swelling over the breasts and chest area.
- ☞ Contact us immediately if you notice significant one-sided swelling or pain.